

AD 679942

OFFICE OF NAVAL RESEARCH


Contract Nonr-4756(07)

Task No. NR 015-816

Determination of the Rate Constants
of the Intersystem Crossing Processes to the
Individual Multiplets of the Lowest Triplet State

M. A. El-Sayed and L. Hall

Department of Chemistry
University of California
Los Angeles, California




Reproduction in whole or in part is permitted for
any purpose of the United States Government

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

October, 1968

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va. 22151



DETERMINATION OF THE RATE CONSTANTS
OF THE INTERSYSTEM CROSSING PROCESSES TO THE
INDIVIDUAL MULTIPLETS OF THE LOWEST TRIPLET STATE

M. A. El-Sayed[†] and L. Hall[‡]

Department of Chemistry*
University of California
Los Angeles, California

The decay of the phosphorescence emission from the three individual sublevels of the lowest triplet state has recently been reported⁽¹⁾ for the first time for pyrazine at 1.6°K. This is observed when the spin-lattice relaxation processes between the three sublevels are slower than the radiative processes from the individual sublevels.^{(1),(2)} Furthermore, the three observed emission lifetimes should be different so that they may be resolved in time. There are a good number of molecules for which these conditions are satisfied.⁽³⁾ It thus appears that one can study the radiative and nonradiative properties of the individual sublevels of the lowest triplet state of phosphorescent molecules.

A number of properties are now being examined in our laboratories. We report here on a new method that can be used to determine the rate constants of the intersystem crossing processes leading to the population of the three different sublevels of the lowest triplet states of phosphorescent molecules. A knowledge of these constants (or their ratios) is vital in determining the relative importance of the different routes of the intersystem crossing processes in molecules.

[†] Alfred P. Sloan Fellow.

[‡] Part of Ph.D. dissertation at UCLA.

* Contribution No. 2342 from UCLA.

The System:

1. A molecule excited to a singlet state loses its energy by internal conversion processes and finally reaches the lowest singlet state, S_1 .
2. Intersystem crossing process proceeds from S_1 to the different triplet sublevels τ_1 , τ_2 and τ_3 of the lowest triplet state, either directly or indirectly via other triplet states located between S_1 and the lowest triplet state. The pumping rates of the τ_1 , τ_2 and τ_3 sublevels from S_1 have the rate constants k_1 , k_2 and k_3 , respectively. Each one of these rate constants may or may not be a composite of other rate constants, depending on the exact mechanism involved in the $S_1 \rightarrow \tau_i$ radiationless transitions.
3. The system is at temperatures for which the S.L.R. processes between τ_1 , τ_2 and τ_3 are much slower than the radiative and nonradiative processes from the three sublevels to the ground state.

Simple Relationships:

The application of the steady state approximation to the formation and disappearance of any of the multiplets, e.g., τ_i , of the above system gives the following relation:

$$K_{(IS)i}[S_1] = K_i[\tau_i] \quad (1)$$

where $K_{(IS)i}$ is the rate constant for the intersystem crossing process from S_1 to the τ_i sublevel, $K_i (=K_{(rad)i} + K_{(nonradiative)i})$ is the observed rate constant of the decay of sublevel i , and $[S_1]$, $[\tau_i]$ represent the steady state concentrations of the lowest singlet state and sublevel (i) of the lowest triplet state, respectively. The observed steady state intensity, I_i , of the emission from sublevel τ_i is given by:

$$I_i = K_{(rad)i}[\tau_i] = K_i[\tau_i]\varphi_i = K_{(IS)i}[S_1]\varphi_i \quad (2)$$

where $\varphi_1 = \frac{K_{(rad)1}}{K_1}$ is the phosphorescence quantum yield of sublevel 1. Applying equation (2) to the emission from any two of the sublevels, e.g., τ_1 and τ_2 , one obtains:

$$\frac{I_1}{I_2} = \frac{K_{(IS)1}}{K_{(IS)2}} \times \frac{\varphi_1}{\varphi_2} \quad (3)$$

If one determines φ_1 , φ_2 and φ_3 at these temperatures, two independent equations can be obtained relating $\frac{I_1}{I_2}$ and $\frac{I_1}{I_3}$ to ratios involving $K_{(IS)1}$. If the total K_{IS} ($= K_{(IS)1} + K_{(IS)2} + K_{(IS)3}$) is determined in the usual manner, one can solve for $K_{(IS)1}$, $K_{(IS)2}$ and $K_{(IS)3}$ individually.

Applications to Pyrazine-d₄:


Since φ_1 , φ_2 and φ_3 are not yet known for any molecule, we selected a molecule with very high phosphorescent quantum yield, pyrazine-d₄. The change of the phosphorescence lifetime upon deuteration is found to be absent for the short-lived emission and only small for the medium- and long-lived emissions. These results strongly suggest that the observed lifetime is near the radiative one for a molecule having as high a triplet-state energy as pyrazine.⁽⁵⁾

These facts justify making the assumption that $\varphi_1 = \varphi_2 = \varphi_3 = 1$. The initial intensity of the three decay components is determined from the decay curve. From polarization and theoretical considerations⁽⁶⁾ the short-, medium-, and long-lived emissions are assigned to radiation from the τ_y , τ_z and τ_x sublevels of the $^3B_{3u}$ state, respectively.[†] Using equation (3), one determines the ratios $\frac{K_{IS}(\tau_z)}{K_{IS}(\tau_y)}$ and $\frac{K_{IS}(\tau_x)}{K_{IS}(\tau_y)}$ to be 2% and 4%, respectively. This means

that the most probable intersystem crossing route in pyrazine is the one(s) that populates the τ_y sublevel of the $^3B_{3u}^{n,\pi^*}$ state. Such a conclusion is in agreement with predictions made previously⁽⁷⁾ which proposed that the most

[†] The N...N and the out-of-plane axes are designated as the z and x axes, respectively.

probable intersystem crossing process in pyrazine is:



The ${}^1B_{3u}^{n,\pi^*} \xrightarrow{s.o.} {}^3B_{1u}^{\pi,\pi^*}$ is the only route allowed by spin-orbit selection rules. The other two levels, τ_z and τ_x , are thus populated by processes involving spin orbit-vibronic or spin-vibronic type interaction. From the determined ratio, it is clear that these processes are 25-50 times less probable than the first-order-allowed processes in pyrazine.

The application of this method to other systems is now in progress, and a complete account of the work will be published later.

Acknowledgements:

We wish to acknowledge the financial support of the Office of Naval Research, Contract No. Nonr-4756(07).

REFERENCES

1. a. L. Hall, A. Armstrong, W. Moomaw, and M. A. El-Sayed, J. Chem. Phys., **48**, 1395 (1968).
b. M. A. El-Sayed, W. R. Moomaw, and D. Tinti, J. Chem. Phys., in press.
c. M. A. El-Sayed, W. R. Moomaw, L. Hall, and A. Armstrong, in International Symposium on Excitons, Magnons and Phonons in Molecular Crystals, A. B. Zahlan, Editor, Cambridge University Press (1968).
2. M. S. de Groot, I. A. Hesselmann, and J. van der Waals, Mol. Phys., **12**, 259 (1967).
3. L. Hall and M. A. El-Sayed, work in progress.
4. a. For review, see: S. K. Lower and M. A. El-Sayed, Chem. Rev., **66**, 199 (1966).
b. D. Scott and O. Maltenkieks, J. Phys. Chem., **72**, 3354 (1968).
5. G. W. Robinson and P. Frosch, J. Chem. Phys., **37**, 1962 (1962); **38**, 1187 (1963).
6. M. A. El-Sayed, "The Intersystem Crossing to and the Phosphorescence from the Individual Sublevels of the Lowest Triplet State in Pyrazine at 1.6°K", in International Conference on Molecular Luminescence (Loyola University, August, 1968), E. Lim, Editor; W. Benjamin, Inc., in press.
7. M. A. El-Sayed, J. Chem. Phys., **36**, 573 (1962); **38**, 2834 (1963).

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) University of California, Los Angeles M. A. El-Sayed, Principal Investigator Los Angeles, California		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP none	
3. REPORT TITLE DETERMINATION OF THE RATE CONSTANTS OF THE INTERSYSTEM CROSSING PROCESSES TO THE INDIVIDUAL MULTIPLETS OF THE LOWEST TRIPLET STATE			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Technical report			
5. AUTHOR(S) (First name, middle initial, last name) M. A. El-Sayed and L. Hall			
6. REPORT DATE October, 1968		7a. TOTAL NO. OF PAGES 5	7b. NO. OF REFS 7
8a. CONTRACT OR GRANT NO. Nonr-4756(07)		9a. ORIGINATOR'S REPORT NUMBER(S)	
b. PROJECT NO. ARPA Order No. 306 Task No. NR 015-816			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.			
10. DISTRIBUTION STATEMENT Distribution of this document is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Office of Naval Research Physics Branch Washington, D. C.	
13. ABSTRACT Equations are derived which can be used to determine the rate constants for the intersystem crossing processes involved in populating the three zero field spin levels of the lowest triplet states of phosphorescing molecules. Such as $\text{P}(\text{P}^1, \text{P}^3, \text{P}^5)$			

14.

KEY WORDS

LINK A

LINK B

LINK C

ROLE

WT

ROLE

WT

ROLE

WT

intersystem crossing
radiationless transitions
pyrazine